

Meta-Analysis of Psychosocial Interventions for Caregivers of People with Dementia

Henry Brodaty, MD, FRANZCP, FRACP, *† Alisa Green, B. Sc (Psychol). Hons,† and Annette Koschera, PhD†

OBJECTIVES: To review published reports of interventions for caregivers (CGs) of persons with dementia, excluding respite care, and provide recommendations to clinicians.

DESIGN: Meta-analytical review. Electronic databases and key articles were searched for controlled trials, preferably randomized, published in English from 1985 to 2001 inclusive. Thirty studies were located and scored according to set criteria, and the interventions' research quality and clinical significance were judged.

SETTING: Home or noninstitutional environment.

PARTICIPANTS: Informal CGs—persons providing unpaid care at home or in a noninstitutional setting.

MEASUREMENTS: The primary measures were psychological morbidity and burden. Other varied outcome measures such as CG coping skills and social support were combined with measures of psychological distress and burden to form a main outcome measure.

RESULTS: The quality of research increased over the 17 years. Results from 30 studies (34 interventions) indicated, at most-current follow-up, significant benefits in caregiver psychological distress (random effect size (ES) = 0.31; 95% confidence interval (CI) = 0.13–0.50), caregiver knowledge (ES = 0.51; CI = 0.05–0.98), any main caregiver outcome measure (ES = 0.32; CI = 0.15–0.48), and patient mood (ES = 0.68; CI = 0.30–1.06), but not caregiver burden (ES = 0.09; CI = –0.09–0.26). There was considerable variability in outcome, partly because of differences in methodology and intervention technique. Elements of successful interventions could be identified. Success was more likely if, in addition to CGs, patients were involved. Four of seven studies indicated delayed nursing home admission.

CONCLUSION: Some CG interventions can reduce CG psychological morbidity and help people with dementia

stay at home longer. Programs that involve the patients and their families and are more intensive and modified to CGs' needs may be more successful. Future research should try to improve clinicians' abilities to prescribe interventions. *J Am Geriatr Soc* 51:000–000, 2003.

Key words: meta-analysis; family caregivers; dementia; interventions

Most people with dementia have at least one supporter or caregiver (CG), usually a spouse or relative. CGs experience adverse psychological, physical, social, and financial consequences,^{1,2} such as higher rates of depression,^{3,4} poorer physical health than non-CG controls,⁵ social isolation,⁶ and direct (e.g., medications) and indirect (e.g., loss of earnings due to relinquishing of paid work) financial costs. CGs are crucial for maintaining people affected with dementia in the community. When there is no CG, or when the CG is stressed, the likelihood of nursing home admission rises sharply.⁷

Clinicians and researchers have devised many methods of trying to help CGs such as education and training programs, support groups, and counseling. Successful interventions have been reported to reduce CG distress, depression, and psychological morbidity; to delay nursing home admission of patients; and to improve patients' psychological well-being. The aim of this study was to review the evidence for the outcome of CG interventions (excluding respite care, which is a patient-targeted intervention—see Gottlieb et al. for a review)⁸ and to provide recommendations for clinicians. The review has been restricted to studies involving informal CGs (persons providing unpaid care, at home or in a noninstitutional environment).

METHOD

Key words (caregiver, carer, self-help groups, support groups, education, training, skills training, counseling, psychotherapy, intervention, and therapy) were used to search published literature in English for controlled studies of interventions for CGs of people with dementia and were each combined with the search terms “random allocation” and/or “control group” and “dementia” or “Alz-

From the *School of Psychiatry, University of New South Wales, Sydney, Australia; and †Academic Department for Old Age Psychiatry, Prince of Wales Hospital, Randwick, New South Wales, Australia

Address correspondence to Professor Henry Brodaty, Academic Department for Old Age Psychiatry, The Euroa Center, Prince of Wales Hospital, Avoca Street, Randwick, Sydney, NSW 2031, Australia. E-mail: h.brodaty@unsw.edu.au

heimer's disease." The following electronic databases were searched: Medline (1985–Week 4, December 2000), PsychInfo (1984–Week 2, December 2000), Ageline (1985–2000/12), CINAL (1985–2000), Cochrane Library 1998 (Issue 3 Database), EBM Reviews—Best Evidence (1991–November/December 2000), EBM Reviews—Cochrane Database of Systematic Reviews (4th Quarter 2000), EMBASE (1988–Week 51, 2000).

All randomized or quasi-experimental trials in which CGs were allocated to intervention or nonintervention (control) groups were selected. Participants in the selected studies were informal CGs of people diagnosed with Alzheimer's disease.

One of the investigators (AG) scrutinized the resulting abstracts, and papers judged relevant were obtained. If doubt existed regarding an article's relevance, a second investigator (HB) reviewed the abstract. References of obtained papers were reviewed to locate any additional papers. Fifty-two articles met criteria for inclusion, seven of which, as shown in brackets in Table 1, were excluded because they were descriptions of earlier studies published by the same research group and added no new information.

The primary outcome measures were psychological morbidity and burden. Measures of psychological morbidity included the General Health Questionnaire, Hamilton Depression Rating Scale, Brief Symptom Inventory, Self-Rating Depression Scale, Hopkins Symptom Checklist, Center for Epidemiological Studies Depression Scale, and the Positive and Negative Affect Scale. Burden was measured using the Burden Interview, Rankin Scale, Caregiver Hassles Scale, Screen for Caregiver Burden, or other objective burden scales.

For those studies that did not include a measure of psychological morbidity or burden, the varied outcome

measures used were combined with measures of psychological distress and burden to form "any main outcome measure." These included measures of CG coping skills (Health Specific Family Coping Index) and social support (Instrumental and Expressive Social Support scale). Knowledge of Alzheimer's disease was examined separately and was measured using the Alzheimer's Disease Knowledge Test, Dementia Quiz, Dementia Knowledge Test, and other individual knowledge measures. "Study success," categorized as a dichotomous dependent variable, was defined as significant change in one of the main outcome measures or an effect size (ES) of 0.5 or greater.

A small number of studies looked at patient mood as an outcome, measured using the Cornell Scale for Depression in Dementia and the Geriatric Depression scale. Finally, the effects of CG interventions on nursing home placements were examined.

Follow-up points were classified as posttest, 3- to 6-month follow-up, more than 6-month follow-up, and most-current follow-up of each study. For the statistical analyses of study characteristics (see below) the most-current follow-up was the dependent variable.

Both reviewers (AG and HB) independently rated the methodological quality of the included studies according to criteria based upon Cochrane Collaboration Guidelines⁹ (Table 2). Characteristics of the design, subjects, outcomes, statistics, and results were used to evaluate the quality of studies (Table 1, quality scores).

Interventions were also rated in terms of their "dosage" or the "strength" of the intervention (Table 1), where the number of sessions/occasions of contact were rated as a minimal (1–2 sessions), moderate (3–5 sessions), medium-high (6–10 sessions), or high/intensive (>10 sessions).

Meta-analysis was performed using MetaView version 4.0 (Cochrane Collaboration, Oxford, England). ES for continuous data was calculated as standardized mean difference (SMD (Cohen's *d*) with 95% confidence intervals (CIs)) between treatment and control group.^{48,50,51} An ES of 0.2 may be statistically significant but is considered weak, 0.5 is considered moderate, and 0.8 or above as strong.⁵² ESs for dichotomous outcome data were reported as odds ratios (ORs).⁵³ Weighted average ESs were calculated weighting each individual ES by the inverse of its variance.⁵³ All pooled calculations included a test of homogeneity of means. Results were compared for fixed-effects and random-effects models.⁵⁴

In most cases, there was no substantial difference between fixed-effects and random-effects models. Results for random-effects models are displayed because tests for homogeneity and heterogeneity of studies under examination in terms of methods, type of intervention, sample characteristics, and outcome measures support the use of a random-effects model for most of the pooled estimates.^{48,49,54} Sensitivity analyses were conducted using various combinations of trials and estimates.

SPSS version 10.0 (SPSS Inc., Chicago, IL) was used to analyze predictors of ES. Chi-square analysis was employed for dichotomous data and Mann-Whitney *U* tests (two-tailed; denoted as *U*) for continuous data. Spearman rank correlation coefficient (Spearman rho) was used as a measure of association.

Where studies contained more than one intervention

Q1

Q2

Q3

Table 1. Criteria for Rating Quality of Studies

Criterion	Score
Design	
Randomized	1
Controlled (or comparison group used)	1
Subjects	
Use of standardized diagnostic criteria	1
All subjects accounted for/withdrawals noted	1
Outcomes	
Well-validated, reliable measures (caregiver and/or patient)	1
Objective outcome (e.g., institutionalization or death)	1
Questionable/unreliable outcome measures	0
Statistics	
Statistical significance considered	1
Adjustment for multiple comparisons	1
Evidence of sufficient power	1
Results	
Blind ratings	1
Follow-up assessment 6 months or beyond	1
Good quality	>7
Poor quality	<5

Table 2. Characteristics of Included Studies

Study	Design	Outcome Measure*	Instrument	Number of Subjects Randomized†	Intervention Type	Quality	Dosage
Brennan et al. 1995 ¹⁰ (same sample as Bass et al. 1998) ¹¹	RCT	Any main outcome measure	IESS	102 (51 each group) Withdrawals = 6	S	6	4
Brodady & Gresham, 1989 ¹² (same sample as Brodady et al. 1991, 1997) ^{13,14}	RCT	Psychological morbidity Delay to nursing home admission†	GHQ	100 (tmt group 1 = 33, tmt group 2, = 31, control = 32)	E, S, C, F, SM, P	7	4
Brodady et al. 1994 ¹⁵	NR	Psychological morbidity Burden Knowledge	GHQ BI Knowledge measure	81 (completed = 33, partially completed = 22, control = 26)	E, S, SM	6	3
Chang et al. 1999 ¹⁶	RCT	Psychological morbidity Burden	BSI CAT	87 (65 completed: tmt = 31, control = 34)	P	6	3
Chiverton & Caine, 1989 ¹⁷	NR	Any main outcome measure Knowledge	HSFCI HSFCI	47 (40 completed: 20 each group)	E, S	5	2
Chu et al. 2000 ¹⁸	RCT	Delay to nursing home admission†		75 (tmt = 37, control = 38) withdrawals = 6	T	7	4
Dröes et al. 1999 ¹⁹	NR	Patient mood	CDS	56 (tmt = 33, control = 23) withdrawals = 14	E, S, P	6	4
Eloniemi-Sulkava et al. 1999 ²⁰	RCT	Delay to nursing home admission†		100 (tmt = 53, control = 47)	E, C, P	7	4
Gendron et al. 1996 ²¹	RCT	Psychological morbidity Burden	HSC BI	35 (tmt = 17, control = 18) withdrawals = 9	S, T	7	3
Hebert et al. 1994 ²² (same sample as Herbert et al. 1995) ²³	RCT	Psychological morbidity Burden Knowledge Delay to nursing home admission†	BSI BI ADKT	45 (tmt = 24, control = 21) withdrawals = 7	S, SM	8	3
Hinchliffe et al. 1995 ²⁴	RCT	Psychological morbidity	GHQ	40 (tmt = 22, control = 18) withdrawals = 14	E, S, C, SM, P	8	4
Kahan et al. 1985 ²⁵	NR	Psychological morbidity Burden Knowledge	SDS BI Dementia Quiz	40 (tmt = 22, control = 18)	S	3	3
LoGiudice et al. 1999 ²⁶	RCT	Psychological morbidity Burden Knowledge	GHQ BI DKT	50 (25 each group) withdrawals = 5	C, F, P	7	1
Marriot et al. 2000 ²⁷	RCT	Psychological morbidity Patient mood	GHQ CSDD	42 (3 groups of 14) withdrawals = 1	E, S, SM, T	9	4

(Continued)

Table 2. (Continued)

Study	Design	Outcome Measure*	Instrument	Number of Subjects Randomized†	Intervention Type	Quality	Dosage
McCallion et al. 1999 ²⁸	RCT	Burden	CHS	66 (tmt = 32, control = 34) withdrawals = 9	E, P	10	3
McCurry et al. 1998 ²⁹	RCT	Patient mood Psychological morbidity Burden	CSDD CES-D SCB	36 (tmt1 = 7, tmt2 = 15, control = 15) withdrawals = 2	S, T, SM	6	3
Mittelman et al. 1996 ³⁰ (same sample as Mittelman et al. 1993, 1995) ^{31,32}	RCT	Delay to nursing home admission†		206 (103 each group) withdrawals = 15	S, C, F	7	4
Mittelman et al. 1995 ³²	RCT	Psychological morbidity	GDS	206 (withdrawals = 9)	S, C, F	7	4
Mohide et al. 1990 ³³ (same sample as Drummond et al. 1991) ³⁴	RCT	Psychological morbidity Delay to nursing home admission†	CES-D	60 (30 each group)	E, S, F	7	4
Moniz-Cook et al. 1998 ³⁵	NR	Psychological morbidity	GHQ	30 (15 each group) withdrawals = 5	P	8	3
Morris et al. 1992 ³⁶	NR	Psychological morbidity Knowledge	BDI Knowledge questionnaire	39 (tmt = 13, control = 18) withdrawals = 8	SM	4	2
Ostwald et al. 1999 ³⁷	RCT	Psychological morbidity Burden	CES-D BI	117 (tmt = 72, control = 45) withdrawals = 23	E, F	5	3
Quayhagen et al. 1989 ³⁸	NR	Psychological morbidity Burden	HSC BI	16 (tmt = 10, control = 6) withdrawals = 4	T, P	6	4
Quayhagen et al. 2000 ³⁹ – Day care – Dyadic counseling – Cognitive stimulation	RCT	Psychological morbidity Burden	BSI MBPC-B	103 (group n's = 21, 29, 22, 16, 15)	C, P, S S, P C P	7	Day care = 4 Dyad counsel = 3 Cog stim = 3
Riordan et al. 1998 ⁴⁰	NR	Delay to nursing home admission†		38 (19 each group) withdrawals = 15	S	5	4
Ripich et al. 1998 ⁴¹	NR	Psychological morbidity Knowledge	PANAS CHS Knowledge questionnaire	37 (tmt = 19, control = 18)	E	5	2
Roberts et al. 1999 ⁴²	RCT	Psychological morbidity	PAIS	77 (tmt = 38, control = 39) withdrawals = 19	C	7	3
Robinson et al. 1988 ⁴³ (same sample as Robinson & Yates, 1994) ⁴⁴	RCT	Burden	Objective burden scale	20 (tmt = 11, control = 9)	S, T	4	2

(Continued)

Table 2. (Continued)

Study	Design	Outcome Measure*	Instrument	Number of Subjects Randomized†	Intervention Type	Quality	Dosage
Teri et al. 1997 ⁴⁵ Problem solving Pleasant events	RCT	Psychological morbidity Burden Patient mood	HDRS BI CDS	88 withdrawals = 16	T, P	7	Problem solving = 3 Pleasant events = 3
Zanetti et al. 1998 ⁴⁶	NR	Psychological morbidity Burden	BSI Rankin scale	23 (tmt = 12, control = 11) withdrawals = 2	S, T	4	3
Zarit et al. 1987 ⁴⁷ Counseling Support group	RCT	Knowledge Psychological morbidity Burden	ADKT BSI BI	184 (group n's = 44, 36, 39) withdrawals = 65	E, F SM S	7	Counseling = 3 Support = 3

* Category of outcome in which this measure was included.

† Intention-to-treat figures.

‡ Time/delay to nursing home admission was a separate analysis, not part of meta-analysis.

ADKT = Alzheimer's Disease Knowledge Test; BDI = Beck Depression Inventory; BI = Burden Interview; BSI = Brief Symptom Inventory; C = counseling of carer; CAT = Caregiver Appraisal Tool; CES-D = Center for Epidemiological Studies—Depression scale; CDS = Cornell Depression Scale; CHS = Caregiving Hassles Scale; CSDD = Cornell Scale for Depression in Dementia; DKT = Dementia Knowledge Test; E = education; F = family counseling/extended family involvement; GHQ = General Health Questionnaire; HDRS = Hamilton Depression Rating Scale; HSC = Hopkins Symptom Checklist; HSFCI = Health Specific Family Coping Index for Non-Institutional Care; IESS = Instrumental and Expressive Social Support Scale; MBPC-B = Memory and Behavior Problem Checklist, Part B; NR = nonrandomized (Quasi-experimental); P = patient involvement; PAIS = Psychosocial Adjustment to Relative's Illness; PANAS = Positive and Negative Affect Scale; RCT = randomized controlled trial; S = support group/program; SCB = Screen for Caregiver Burden; SDS = Self-Rating Depression Scale; SM = stress management; T = training; tmt = treatment.

group (e.g., support group compared with counseling group vs control), each intervention group was entered into the analysis separately. The following studies were excluded: (1) two studies^{55,56} with a sample size of five or fewer in treatment or control group, (2) 11 studies⁵⁷⁻⁶⁷ with insufficient outcome information to calculate ES or nursing home delay, and (3) two interventions with extreme values (values more than three times the interquartile range; dual seminar intervention (SMD for Brief System Inventory depression = -1.86);³⁹ the other three interventions for this study were included in the analysis), and Perkins et al. 1990⁶⁸ (SMD for caregiver morale = 3.09).⁵¹ Therefore, of the 45 studies that met criteria for inclusion, 30 (34 interventions) were included in at least one analysis.

RESULTS

The 30 controlled studies involved 2,040 CGs (intention-to-treat; range 16-206, median = 53), who were predominantly spouses (of persons with dementia), female, and aged 55 and older (see Table 1 for included studies).

Quality ratings, which ranged from 3 to 10 (mean ± standard deviation = 6.4 ± 1.5), tended to improve over time (Spearman rho = 0.3; P = .07). There was no significant correlation of quality of research on ES (any outcome, distress, burden) using nonparametric measures (Spearman rho = 0.2, 0.3, and 0.2, respectively).

The ESs for psychological morbidity are shown in Table 3. Although 77% of the studies showed a positive ES for psychological morbidity (range -0.59-1.81), this was only statistically significant in five of the 20 positive interventions. The ES for burden ranged from -0.6 (95% CI = -1.33-0.14) to 1.07 (95% CI = 0.12-2.03). Only one intervention of 20 (a social skills training program)⁴³ showed a statistically significant effect on burden. Overall, 23 of 34 (68%) interventions met the criteria for study success.

Weighted average ESs (95% CI; number of studies) were calculated for CG psychological morbidity, 0.31 (0.13-0.50; n = 26); CG burden, 0.09 (-0.09-0.26; n = 20); changes in patient mood, 0.68 (0.30-1.06; n = 5); CG knowledge, 0.51 (0.05-0.98; n = 8); and overall effect on "any main outcome measure," 0.32 (0.15-0.48; n = 30). The weight used here was the inverse of its variance (i.e., larger studies are given more weight).^{53,69} Low but significant ESs on most outcome measures (apart from burden) suggest a low but positive overall effect of these CG interventions, but the variability between studies is substantial. The pooled estimates displayed were calculated for the most-current follow-up assessment, which was posttest for most studies. Eight studies reported results for additional follow-up assessments (mean = 27 weeks; range = 12-48 weeks). Pooled estimates for time intervals of 3 to 6 months and more than 6 months are available upon request.

Sensitivity analyses for the most-current assessment point were performed, excluding extreme values (see above), dropping one study at a time and obtaining CIs for the remaining studies.⁵¹ Certain studies proved more influential than others, yet discrepancies in CIs for pooled calculations were small and did not result in a change of significance when compared with the overall result. An

Table 3. Effect Size for Psychological Morbidity at Most Current Follow-Up Assessment

Study	Standardized Mean Difference* (95% Confidence Interval)
Moniz-Cook et al. 1998 (GHQ)	1.81 (0.94–2.67)
Marriot et al. 2000 (GHQ)	1.57 (0.69–2.45)
Hinchliffe et al. 1995 (GHQ)	1.42 (0.64–2.21)
Teri et al. 1997; problem solving (HDRS)	1.10 (0.27–1.92)
Quayhagen et al. 1989 (HSC)	0.92 (–0.16–2.00)
Brodaty and Gresham 1989 (GHQ)	0.77 (0.27–1.28)
Quayhagen et al. 2000; cog. stimulation (BSI)	0.59 (–0.09–1.27)
Teri et al. 1997; pleasant events (HDRS)	0.53 (–0.23–1.29)
Zanetti et al. 1998 (BSI)	0.46 (–0.42–1.34)
Chang et al. 1999 (BSI)	0.45 (–0.04–0.95)
Mittelman et al. 1995 (GDS)	0.29 (0.02–0.60)
Mohide et al. 1990 (CES-D)	0.26 (–0.35–0.87)
Ostwald et al. 1999 (CES-D)	0.25 (–0.20–0.70)
McCurry et al. 1998 (CES-D)	0.21 (–0.58–1.00)
Hebert et al. 1994 (BSI)	0.20 (–0.47–0.86)
Ripich et al. 1998 (PANAS)	0.15 (–0.50–0.81)
Quayhagen et al. 2000; day care (BSI)	0.12 (–0.58–0.83)
Kahan et al. 1985 (SDS)	0.09 (–0.53–0.72)
Gendron et al. 1996 (HSC)	0.07 (–0.60–0.73)
Zarit et al. 1987; counseling (BSI)	0.02 (–0.43–0.48)
Morris et al. 1992 (BDI)	–0.09 (–0.80–0.63)
Brodaty et al. 1994 (GHQ)	–0.16 (–0.71–0.38)
Zarit et al. 1987; support group (BSI)	–0.17 (–0.60–0.27)
Logiudice et al. 1999 (GHQ)	–0.18 (–0.87–0.52)
Roberts et al. 1999 (PAIS)	–0.24 (–0.75–0.28)
Quayhagen et al. 2000; dyadic counseling (BSI)	–0.59 (–1.23–0.05)

* Effect size measured as standardized mean difference between treatment and control group.

GHQ = General Health Questionnaire (Goldberg & Williams, 1988); BSI = brief symptom inventory (Derogatis et al. 1983); SDS = Self-rating Depression Scale (Zung, 1965); HSC = Hopkins Symptom Checklist (Derogatis et al. 1974); CES-D = Center for Epidemiological Studies Depression Scale (Radloff, 1977); PANAS = Positive and Negative Affect Scale (modified version; Watson et al. 1988); PAIS = Psychosocial Adjustment to Illness Scale.

exception was the sensitivity analysis for knowledge, for which the overall SMD for random effect models was significant, whereas removal of any of the more successful studies led to nonsignificant results.

A post hoc analysis of study characteristics was used to test possible predictors of positive ES. The following predictor variables were examined: whether the intervention involved support/help from extended family; counseling of the CG; and involvement of CG and patient in intervention (e.g., teaching the caregiver skills applicable to the patient such as pleasant event planning, cognitive stimulation), support group, and stress management. Continuous dependent variables were the ES (SMD for random effect models) for CG psychological morbidity and the ES on any main outcome measure.

Using univariate analyses, the study characteristic “involvement of CG and patient in the intervention” showed an effect on any outcome measure ($U = 43.0$; $P = .01$), on CG psychological morbidity ($U = 28$; $P = .01$) and on study success ($\chi^2 = 4.0$; $df = 1$; continuity correction: $P = .05$). No other study characteristic showed a significant effect.

The dosage of interventions ranged from minimal ($n = 1$), to moderate ($n = 4$), medium-high ($n = 17$), and high/intensive ($n = 12$). Higher dosage was associated with de-

creased psychological distress (Spearman rho = 0.6; $P = .003$), but there was a nonlinear tendency with interventions with a dosage of greater than 3.5 being less effective than interventions around 3.5.

Seven studies used time until nursing home placement as an outcome measure.^{12,18,20,22,30,33,40} Two of these showed significant ESs (Brodaty et al.¹² OR = 5.0, 95% CI = 1.72–14.70; Mittelman et al.³⁰ SMD = 3.21, 95% CI = 2.80–3.63), and two reported a longer median time of home care until institutionalization in the intervention group than in the control group (Eloniemi-Sulkava et al.²⁰ 2-year follow-up: median time of home care in those patients who were institutionalized 473 vs 240 days, respectively; $P = .02$; Riordan et al.⁴⁰ 326 vs 160 days, respectively). Chu et al.¹⁸ divided patients into very mildly and mildly to moderately impaired. Those in the latter class who received treatment remained in the community an average of 52.53 days longer than control group patients. Delays in nursing home admission of between 53 and 329 days were reported.^{12,18,20,30,40} Two studies^{35,40} reported that significantly more control patients received permanent residential care at follow-up assessment. Qualitatively, a continuing relationship between helper and CG, flexibility of the intervention, and a variety of interven-

tions that should meet the varied needs of CGs appeared to characterize the four studies that delayed nursing home admission.^{12,20,30,40}

DISCUSSION

CG interventions have modest but significant benefits on CG knowledge, psychological morbidity, and other main outcome measures (such as coping skills and social support). At the most-current follow-up, there was a mean ES of 0.3 for all CG outcomes and for CG psychological morbidity in particular, meaning that the average patient in the treatment group was less depressed than about 62% of patients in the control group. There was an even stronger ES (0.7 for posttest and 0.5 for most-current follow-up) in the increase in CG knowledge (about dementia and how to cope with it). There was also a strong ES for patient mood, but interventions did not appear to influence CG burden.

The findings regarding the predictors of positive ES are based on small numbers and should be interpreted with caution. The heterogeneity of sample characteristics and study design contribute a considerable amount of variance but cannot be controlled because of lack of information and the small number of studies. Additionally, the power of the analyses to detect small to medium ESs was less than 0.5 ($\alpha = .05$).⁵²

Despite these modest findings, CGs were frequently satisfied or very satisfied with their interventions,^{15,21,22,25,36,40,42} appraised their own coping skills as improved,¹⁷ reported that their relationship with the patient had improved,²⁸ identified helpful training elements,³⁹ and mostly (71%) reported that they would use training again.⁴³ CG interventions can have effects on delaying nursing home admission, which for many is desirable. Successful interventions are short educational programs (beyond enhancement of knowledge);^{14,27} support groups alone, single interviews, and brief interventions or courses that were not supplemented with long-term contact do not work.

The variability in outcomes is attributable to many factors. Patients were heterogeneous with regard to age, sex, and living arrangements; type and severity of dementia; and prevalence of behavioral and psychological symptoms associated with dementia. CGs also differed significantly with regard to demographic variables, relationship to patient (e.g., spouse vs adult child vs other), other demands on their time (working, other family members to care for), and practical and social supports. The different methods of recruitment (volunteers, clinics, advertisements, and Alzheimer's associations) may have introduced bias. Whether the different follow-up periods influenced the results was considered. Eleven studies had more than one posttest assessment. In these studies, the average ES for psychological distress improved, whereas the ES for burden decreased, but these findings would have to be corrected for many covariates and could not be considered representative of all 30 studies. Finally, the number of subjects in trials was small; there was limited power, statistical comparisons were multiple (corrections for these were few), and intention-to-treat analyses were largely not performed.

Despite these limitations, it is clear that some inter-

ventions can make a difference. What are the important elements? Statistically, the only feature that emerged as significant was involvement of the patient in addition to the CG in a structured program, such as teaching the CG problem-solving skills in the care of the patient.⁴⁵ The small numbers of subjects in a large number of trials may have militated against the emergence of other features that appear qualitatively important: practical support for the CG, involvement of the extended family, structured individual counseling, and a flexible provision of a consistent professional to provide long-term support. Ceiling and floor effects prevented the realization of beneficial effects. For example if only a minority of CGs were significantly depressed before intervention, this limited the possibility of demonstrating a significant reduction in depression score. The difficulty in recruiting and treating sufficient numbers of subjects precluded the examination of interactions between which intervention for which CG supporting which patient with what dementia at what time in the course of the condition.

The implications from this meta-analysis are that CG interventions have the potential to benefit patients and CGs. The quality of research is advancing, but there is considerable room for methodological improvement. Future research should be conducted with more rigor: randomized, controlled, blind outcome assessments, follow-ups for at least 6 months, and use of well-validated and reliable outcome criteria measuring outcomes proximally (burden, knowledge) and distally (depression, quality of life). The next steps include evaluation of more-intensive interventions and interactions with drug therapies.

ACKNOWLEDGMENTS

Many thanks to Dusan Hadzi-Pavlovic for invaluable statistical advice.

REFERENCES

1. Max W, Webber PA, Fox PJ. Alzheimer's disease. The unpaid burden of caring. *J Aging Health* 1995;7:179-199.
2. Brodaty H, Green A. Family caregivers for people with dementia. In: O'Brien J, Ames D, Burns A, eds. *Dementia*, 2nd Ed. London: Arnold, 2000, pp. 193-205.
3. Baumgarten M, Battista RN, Infante-Rivard C et al. The psychological and physical health of family members caring for an elderly person with dementia. *J Clin Epidemiol* 1992;45:61-70.
4. Rosenthal CJ, Sulman J, Marshall VW. Depressive symptoms in family caregivers of long-stay patients. *Gerontologist* 1993;33:249-257.
5. Schulz R, Vistainer P, Williamson GM. Psychiatric and physical morbidity effects of caregiving. *J Gerontol* 1990;45:P181-P191.
6. Brodaty H, Hadzi-Pavlovic D. Psychosocial effects on carers of living with persons with dementia. *Aust N Z J Psychiatry* 1990;24:351-361.
7. Brodaty H, McGilchrist C, Harris L et al. Time until institutionalization and death in patients with dementia: Role of caregiver training and risk factors. *Arch Neurol* 1993;50:643-650.
8. Gottlieb BH, Johnson J. Respite programs for caregivers of persons with dementia: A review with practice implications. *Aging Ment Health* 2000;4:119-129.
9. Clarke M, Oxman AD, eds. *Cochrane Reviewers Handbook 4.1.1* In: *The Cochrane Library*, Oxford: Update Software. 2000.
10. Brennan FP, Moore SM, Smyth KA. The effects of a special computer network on caregivers of persons with Alzheimer's disease. *Nurs Res* 1995;44:166-172.
11. Bass DM, McClendon MJ, Flatley Brennan P et al. The buffering effect of a computer support network on caregiver strain. *J Aging Health* 1998;10:20-43.
12. Brodaty H, Gresham M. Effect of a training programme to reduce stress in carers of patients with dementia. *BMJ* 1989;299:1375-1379.
13. Brodaty H, Gresham M, Luscombe G. The Prince Henry Hospital dementia caregivers training programme. *Int J Geriatr Psychiatry* 1997;12:183-192.

14. Brodaty H, Peters KE. Cost effectiveness of a training program for dementia carers. *Int Psychogeriatr* 1991;3:11–21.
15. Brodaty H, Roberts K, Peters K. Quasi-experimental evaluation of an educational model for dementia caregivers. *Int J Geriatr Psychiatry* 1994;9:195–204.
16. Chang BL. Cognitive-behavioural intervention for homebound caregivers of persons with dementia. *Nurs Res* 1999;48:173–182.
17. Chiverton P, Caine ED. Education to assist spouses in coping with Alzheimer's disease. A controlled trial. *J Am Geriatr Soc* 1989;37:593–598.
18. Chu P, Edwards J, Levin R et al. The use of clinical case management for early stage Alzheimer's patients and their families. *Am J Alzheimers Dis Other Demen* 2000;15:284–290.
19. Dröes RM, Breebaart E, Ettema TP et al. Effect of integrated family support versus day care only on behaviour and mood of patients with dementia. *Int Psychogeriatr* 1999;12:99–115.
20. Eloniemi-Sulkava U, Sivenius J et al. Support program for demented patients and their carers: The role of dementia family care coordinator is crucial. In: Iqbal K, Swaab DF, Winblad B et al, eds. *Alzheimer's Disease and Related Disorders*. West Sussex: John Wiley & Sons, 1999, pp. 795–802.
21. Gendron C, Poitras L, Dastoor DP et al. Cognitive-behavioral group intervention for spousal caregivers. Findings and clinical considerations. *Clin Gerontol* 1996;17:3–19.
22. Hébert R, Leclerc G, Bravo G et al. Efficacy of a support programme for caregivers of demented patients in the community: A randomized controlled trial. *Arch Gerontol Geriatr* 1994;18:1–14.
23. Hébert R, Girouard D, Leclerc G et al. The impact of a support programme for care-givers on the institutionalisation of demented patients. *Arch Gerontol Geriatr* 1995;20:129–134.
24. Hincliffe AC, Hyman IL, Blizard B et al. Behavioural complications of dementia—Can they be treated? *Int J Geriatr Psychiatry* 1995;10:839–847.
25. Kahan J, Kemp B, Staples FR et al. Decreasing the burden in families caring for a relative with a dementing illness: A controlled study. *J Am Geriatr Soc* 1985;33:664–670.
26. LoGiudice D, Waltrowicz W, Brown K et al. Do memory clinics improve the quality of life of carers? A randomized pilot trial. *Int J Geriatr Psychiatry* 1999;14:626–632.
27. Marriott A, Donaldson C, Tarrier N et al. Effectiveness of cognitive-behavioural family intervention in reducing the burden of care in carers of patients with Alzheimer's disease. *Br J Psychiatr* 2000;176:557–562.
28. McCallion P, Toseland RW, Freeman K. An evaluation of a family visit education program. *J Am Geriatr Soc* 1999;47:203–214.
29. McCurry SM, Logsdon RG, Vitiello M et al. Successful behavioural treatment for reported sleep problems in elderly caregivers of dementia patients: A controlled study. *J Gerontol B Psychol Sci Soc Sci* 1998;53B:P122–P129.
30. Mittelman MS, Ferris SH, Shulman E et al. A family intervention to delay nursing home placements of patients with Alzheimer's disease. A randomized controlled trial. *JAMA* 1996;276:1725–1731.
31. Mittelman MS, Ferris SH, Shulman E et al. A comprehensive support program. Effect on depression in spouse-caregivers of AD patients. *Gerontologist* 1995;35:792–802.
32. Mittelman MS, Ferris SH, Steinberg G et al. An intervention that delays institutionalisation of Alzheimer's disease patients: Treatment of spouse-caregivers. *Gerontologist* 1993;33:730–740.
33. Mohide EA, Pringle DM, Streiner DL et al. A randomized trial of family caregiver support in the home management of dementia. *J Am Geriatr Soc* 1990;38:446–454.
34. Drummond M, Mohide EA, Tew M et al. Economic evaluation of a support programme for caregivers of demented elderly. *Int J Technol Assess Health Care* 1991;7:209–219.
35. Moniz Cook E, Agar S, Gibson GD et al. A preliminary study of the effects of early intervention with people with dementia and their families in a memory clinic. *Aging Ment Health* 1998;2:199–211.
36. Morris RG, Woods RT, Davies KS et al. The use of a coping strategy focused support group for carers of dementia sufferers. *Counselling Psychol Q* 1992;5:337–348.
37. Ostwald SK, Hepburn KW, Caron W et al. Reducing caregiver burden. A randomized psychoeducational intervention for caregivers of persons with dementia. *Gerontologist* 1999;39:299–309.
38. Quayhagen MP, Quayhagen M. Differential effects of family-based strategies on Alzheimer's disease. *Gerontologist* 1989;29:150–155.
39. Quayhagen MP, Quayhagen M, Corbeil RR et al. Coping with dementia. Evaluation of four nonpharmacologic interventions. *Int Psychogeriatr* 2000;12:249–265.
40. Riordan J, Bennett A. An evaluation of an augmented domiciliary service to older people with dementia and their carers. *Aging Ment Health* 1998;2:137–143.
41. Ripich DN, Zioli E, Lee MM. Longitudinal effects of communication training on caregivers of persons with Alzheimer's disease. *Clin Gerontologist* 1998;19:37–53.
42. Roberts J, Browne G, Milne C et al. Problem-solving counseling for caregivers of the cognitively impaired: Effective for whom? *Nurs Res* 1999;48:162–172.
43. Robinson KM. A social skills training program for adult caregivers. *Adv Nurs Sci* 1988;10:59–72.
44. Robinson K, Yates K. Effects of two caregiver-training programs on burden and attitude toward help. *Arch Psychiatr Nurs* 1994;8:312–319.
45. Teri L, Logsdon RG, Uomoto J et al. Behavioral treatment of depression in dementia patients: A controlled clinical trial. *J Gerontol B Psychol Sci Soc Sci* 1997;52B:P159–P166.
46. Zanetti O, Metitieri T, Bianchetti A et al. Effectiveness of an educational program for demented person's relatives. *Arch Gerontol Geriatr Suppl* 1998;6:531–538.
47. Zarit SH, Anthony CR, Boutselis M. Interventions with care givers of dementia patients: Comparison of two approaches. *Psychol Aging* 1987;2:225–232.
48. Normand SL. Tutorial in biostatistics. Meta-analysis. Formulating, evaluating, combining, and reporting. *Stat Med* 1999;18:321–359.
49. Engels EA, Schmid CH, Terrin N et al. Heterogeneity and statistical significance in meta-analysis: An empirical study of 125 meta-analyses. *Stat Med* 2000;19:1707–1728.
50. Hunter JE, Schmidt FL. *Methods of Meta-Analysis. Correcting Error and Bias in Research Findings*. London: Sage, 1990.
51. Olkin I. Diagnostic statistical procedures in medical meta-analyses. *Stat Med* 1999;18:2331–2341.
52. Cohen J. *Statistical Power Analysis for the Behavioural Sciences*. London: Lawrence Erlbaum, 1988.
53. Clarke M, Oxman AD, eds. *Cochrane Reviewers' Handbook 4.0. Review Manager, Version 4.0*. Oxford: The Cochrane Collaboration, 1999.
54. Hardy R, Thompson SG. Detecting and describing heterogeneity in meta-analysis. *Stat Med* 1998;17:841–856.
55. Gendron CE, Poitras LR, Engels ML et al. Skills training with supporters of the demented. *J Am Geriatr Soc* 1986;34:875–880.
56. Sutcliffe C, Larner S. Counselling carers of the elderly at home: A preliminary study. *Br J Clin Psychol* 1988;27:177–178.
57. Haley WE, Brown L, Levine EG. Experimental evaluation of the effectiveness of group intervention for dementia caregivers. *Gerontologist* 1987;27:376–382.
58. Baldwin BA, Kleeman KM, Stevens GL et al. Family caregiver stress. Clinical assessment and management. *Int Psychogeriatr* 1989;1:185–194.
59. Dellasega C. Coping with care-giving. Stress management for caregivers of the elderly. *J Psychosoc Nurs Ment Health Serv* 1990;28:15–22.
60. Goodman CC, Pynoos J. A model telephone information and support program for caregivers of Alzheimer's patients. *Gerontologist* 1990;30:399–404.
61. Seltzer MM, Litchfield LC, Kaput LR et al. Professional and family collaboration in case management: A hospital-based replication of a community-based study. *Soc Work Health Care* 1992;17:1–22.
62. Weinberger M, Gold DT, Divine GW et al. Social service interventions for caregivers of patients with dementia: Impact on health care utilization and expenditures. *J Am Geriatr Soc* 1993;41:153–156.
63. Farran CJ, Keane-Hagarty E. Multi-modal intervention strategies for caregivers of persons with dementia. In: Light E, ed. *Stress Effects on Family Caregivers of Alzheimer's Patient's: Research and Interventions*. New York: Springer Publishing, 1994, pp. 242–259.
64. Challis D, Von Abendorff R, Brown P et al. Care management and dementia: An evaluation of the Lewisham Intensive Case Management Scheme. In: Hunter S, ed. *Dementia: Challenges and New Directions*. London: Jessica Kingsley, 1997, pp. 139–164.
65. Burgener SC, Bakas T, Murray C et al. Effective caregiving approaches for patients with Alzheimer's disease. *Geriatr Nurs* 1998;19:121–126.
66. Buckwalter KC, Gerdner L, Kohout F et al. A nursing intervention to decrease depression in family caregivers of persons with dementia. *Arch Psychiatr Nurs* 1999;13:80–88.
67. Corbeil RR, Quayhagen MP, Quayhagen M. Intervention effects of dementia caregiving interaction. A stress-adaptation modeling approach. *J Aging Health* 1999;11:9–95.
68. Perkins RE, Poynton CF. Group counselling for relatives of hospitalized presenile dementia patients: A controlled study. *Br J Clin Psychol* 1990;29:287–295.
69. Laird NM, Mosteller F. Some statistical methods for combining experimental results. *Int J Tech Assess Health Care* 1990;6:5–30.

QUERIES

Brodaty et al., #51210

Q1: AU, no brackets in Table 1?

Q2: Do you mean Table 2 here?

Q3: Do you mean Table 2 here?

Q4: AU, set as meant (Table 3)?

Proofs Only